

Detection of fungal pathogens on groundnut seeds in different tribal localities of Dangs district under South Gujarat

KOTGIRE GANESH AND B. P. MEHTA

Department of Plant Pathology, N. M. College Agriculture, Navsari Agricultural University, Navsari, Gujarat

Received : 29.07.2013

Accepted : 18.12.2013

Published : 28.04.2014

Fungal detection on seeds from different locations on different varieties of groundnut was carried out by standard blotter method. Frequency of *A. flavus* at Waghai, Bhenskatri, Kudkas and Gaurya was observed maximum except Pimpri i.e. 40.00, 50.00, 53.00 and 48.00 per cent respectively. Assessment of fungi on moldy and healthy seeds showed total five fungal species were associated with surface sterilized healthy seeds and six fungal species were associated with surface sterilized moldy seeds of Local, GG-2, Junagadh-II and TAG-37 varieties. Frequency of *A. flavus* and *A. niger* were maximum in healthy and moldy seeds of all groundnut varieties.

Key words: Frequency, moldy seeds, varieties, etc.

INTRODUCTION

In Gujarat, groundnut is an important crop and its cultivation increased in tribal area due to increase in biodiversity. The total area under cultivation of groundnut is about 17,954 hectares with the production of 24,560 million tonnes of seeds with average productivity of 1368 kg ha⁻¹. In Dangs, groundnut occupies about 46 ha area with a total production of 6300 million tones with average productivity of 1369 kg ha⁻¹ (Anon., 2012). Groundnut seed is infected with large number of field and storage fungi which are responsible for seed rot, seedling blight, stem rot and foliar infection which subsequently hamper seed health. Deterioration of groundnut seeds due to fungal activity is normally associated with the production of off-colours and flavours, rancidity, discolouration, effects on yield and quality of oil, loss of seed viability and formation of mycotoxins (Twiddy, 1994). Agroclimatic conditions play an important role in infecting the seeds by the fungi at

the time of crop maturity as well as during threshing and further processing of the produce. An attempt has been made to find out the effect of locality, in which crop is grown in the association of important fungal pathogens on seeds of groundnut.

MATERIALS AND METHODS

Collection of seed samples

The peoples in tribal area in Dangs district of south Gujarat cultivated and stored the groundnut seeds in bean for further use. Infected and healthy seed samples of widely cultivated groundnut varieties like Local, GG-2, Junagarh-1 and Tag-37 were collected in brown paper bags from farmer's fields of Waghai, Bhenskatri, Kudkas, Gaurya and Pimpri villages of Dangs district.

Detection and isolation of seed borne fungi

Assessment of fungi associated with groundnut seeds of different locations and moldy and healthy

seeds was carried out from randomly taken 100 seeds of groundnut from each location (Table 1)

Table 1 : Collection of seed samples from farmers' fields of Dangs districts

Village	Variety
Kudkas	Local, GG-2, GJ-11
Bhenskatri	Local, TAG - 37
Waghai	Local, GJ-11, GG-2, TAG-37
Gaurya	Local, GG-2
Pimpri	Local, Tag-37

with three replications by standard blotter method (Bhale *et al.*, 2001). Ten seeds after surface sterilization by 1 per cent sodium hypochlorite solution for one minute, were placed at equi distance on three layers of properly moistened sterilized blotters, and Petri plates were incubated under 12/12 h alternating light and dark period at $25 \pm 2^\circ\text{C}$. Developing fungal growth on each of the piece of seeds of groundnut after seven days was observed under stereoscopic binocular microscope and results were recorded accordingly.

Identification and purification

Various seed-infecting fungi, developed on groundnut seeds, were cultured separately on PDA Petri plates. Each fungal growth was critically observed under microscope for cultural and morphological characteristics described in earlier literature. The same pure culture grown in slants were sent to Indian Type Culture Collection (I.T.C.C.), Division of Mycology and Plant Pathology, Indian Agricultural Research Institute (I.A.R.I), New Delhi for identification and confirmation of isolated fungus. Cultures were maintained on PDA slants for purification by sub-culturing and stored at 5°C for further study. Observations on the fungul flora, associated with the seeds, were recorded after 7 days and per cent frequently of each fungus was calculated. Mean frequently percentage of important fungi were compared and presented in Tables 2 and 3.

RESULTS AND DISCUSSION

Identification of pathogens

The different cultures obtained from groundnut seeds were purified by single spore and hyphal tip isolation technique. The identification of the isolated fungus was done by studying the cultural, morphological characters and by microscopic examination of each of isolate and the fungi

identified as *Sclerotium rolfsii*, *Helminthosporium* sp., *Fusarium equiseti*, *Fusarium moniliforme*, *Aspergillus niger* and *Aspergillus flavus*.

Assessment of fungi on farmers stored seeds of groundnut

The data pertaining to the frequency of fungi on farmers saved seeds of groundnut in their own places were presented in Table 2, revealed that the frequency of fungi varied in all the locations which was in the range of 12.00-40.00, 10.00-50.00, 9.00-53.00, 6.00-48.00 and 6.00-43.00 per cent at Waghai, Bhenskatri, Kudkas, Gaurya and Pimpri villages of Dangs districts respectively.

Frequency of *A. flavus* was observed maximum at all the locations except Pimpri i.e. 40.00, 50.00, 53.00 and 48.00 per cent respectively, whereas, seed samples of pimpri recorded highest frequency of *A. niger* i.e. 46.00 per cent. At Waghai, frequency of *A. niger* (28.00%) was next followed by *F. moniliforme* (22.00%), *Helminthosporium* sp. (16.00%) and *S. rolfsii* (12.00%). Seed samples of Bhenskatri recorded minimum frequency of *Helminthosporium* sp. (10.00%) followed by *S. rolfsii* (14.00%) and *F. equiseti* (16.00%). At Kudkas, frequency of *F. equiseti* was observed

Table 2 : Frequency of fungi in per cent on farmers stored seeds of groundnut in Dangs

Fungi	Locations*				
	Waghai	Bhenskatri	Kudkas	Gaurya	Pimpri
<i>Sclerotium rolfsii</i>	12.00	14.00	10.00	6.00	17.00
<i>Helminthosporium</i> sp.	16.00	10.00	19.00	10.00	22.00
<i>Fusarium equiseti</i>	19.00	16.00	9.00	15.00	6.00
<i>Fusarium moniliforme</i>	22.00	29.00	32.00	24.00	18.00
<i>Aspergillus niger</i>	28.00	38.00	37.00	32.00	46.00
<i>Aspergillus flavus</i>	40.00	50.00	53.00	48.00	43.00

*Average of three replications

lowest i.e. only 9.00 per cent followed by *S. rolfsii* (10.00%) and *Helminthosporium* sp. (19.00%), while, *A. niger* was second highest fungus which recorded 37.00 per cent frequency on seeds. The frequency of *S. rolfsii* at Gaurya and Pimpri was 6.00 and 17.00 per cent respectively and *F. moniliforme* 24.00 and 18.00 per cent respectively. The seed samples of Pimpri recorded minimum frequency of *F. equiseti* (16.00%) and highest frequency of *A. niger* (46.000%). This difference may be attributed to the effect of different environmental conditions occurring in different locations which influenced specific fungal infection.

Table 2 : Frequency of fungi in per cent on healthy and moldy seeds of different groundnut varieties

Fungi	Local*		GG 2*		Junagadh 11*		TAG 37*	
	H	M	H	M	H	M	H	M
<i>Sclerotium rolfsii</i>	2.00	8.00	4.00	11.00	3.00	6.00	3.00	9.00
<i>Helminthosporium</i> sp.	0.00	7.00	7.00	15.00	5.00	13.00	4.00	12.00
<i>Fusarium equiseti</i>	7.00	15.00	0.00	5.00	6.00	11.00	7.00	11.00
<i>Fusarium moniliforme</i>	10.00	17.00	10.00	16.00	5.00	15.00	8.00	14.00
<i>Aspergillus niger</i>	13.00	20.00	14.00	18.00	8.00	16.00	11.00	16.00
<i>Aspergillus flavus</i>	15.00	23.00	17.00	24.00	14.00	26.00	14.00	18.00

*Average of three replications

H-Healthy, M-Moldy

This result tallies with results obtained by Agarwal and Gupta (1989). They reported variation of fungal load observed in location to location. The minimum fungal load was detected on seeds of soybean from Junagadh (3.3%) followed by Akola (5.1%) while it was maximum on seeds from Parbhani (9.4%) which was closely followed by Navsari (8.4%).

Assessment of fungi on moldy and healthy seeds of different groundnut varieties

Results on assessment of fungi associated with different varieties of moldy and healthy seeds of groundnut, carried out by blotter methods, are presented in (Table 3). Overall, five fungal species were associated with surface sterilized healthy seeds of all four varieties and six fungal species were associated with surface sterilized moldy seeds of all varieties.

Frequency of fungi on local variety of healthy seeds recorded *S. rolfsii* (2.00%), *F. equiseti* (7.00%), *F. moniliforme* (10.00%), *A. niger* (13.00%) and *A. flavus* (15.00%), while, on moldy seeds, *S. rolfsii* (8.00%), *Helminthosporium* sp. (7.00%), *F. equiseti* (15.00%), *F. moniliforme* (17.00%), *A. niger* (20.00%) and *A. flavus* (23.00%) were found on seeds.

Similarly, fungal frequency on healthy seeds of GG 2 variety was *S. rolfsii* (4.00%), *Helminthosporium* sp. (7.00%), *F. moniliforme* (10.00%), *A. niger* (14.00%) and *A. flavus* (17.00%) and on moldy seeds of GG-2, the frequency was *S. rolfsii* (11.00%), *Helminthosporium* sp. (15.00%), *F. equiseti* (5.00%), *F. moniliforme* (16.00%), *A. niger* (13.00%) and *A. flavus* (24.00%).

Frequency per cent of fungi associated with healthy seeds on variety of Junagadh-11 and TAG-37 were *S. rolfsii*, *Helminthosporium* sp., *F. equiseti*, *F.*

moniliforme, *A. niger* and *A. flavus* i.e. 3.00 and 3.00, 5.00 and 4.00, 6.00 and 7.00, 5.00 and 8.00, 8.00 and 11.00 and 14.00 and 14.00 per cent respectively, and frequency on moldy seeds were 6.00 and 9.00, 13.00 and 12.00, 11.00 and 11.00, 15.00 and 14.00, 16.00 and 16.00 and 26.00 and 18 per cent respectively.

This result tallies with the result obtained by Kushi and Khare (1979). They observed that lesser number of fungi were associated with apparently healthy seeds whereas, moldy seeds recorded more no. of fungi in sesamum. Thus by and large, healthy and moldy seeds revealed preferential trend of fungal association. Thus it can be found that various fungi induced specific grain abnormality and hampered seed health.

Frequency of fungi on farmers stored seeds of groundnut varied in all the locations which was in the range of 12.00-40.00, 10.00-50.00, 9.00-53.00, 6.00-48.00 and 6.00-43.00 per cent at Waghai, Bhenskatri, Kudkas, Gaurya and Pimpri villages of Dangs districts respectively. Frequency of *A. flavus* was observed maximum at all the locations except Pimpri i.e. 40.00, 50.00, 53.00 and 48.00 per cent respectively. Five fungal species were associated with surface sterilized healthy seeds and six fungal species were associated with surface sterilized moldy seeds of Local, GG-2, Junagadh 11 and TAG 37 varieties. Frequency of *A. flavus* and *A. niger* was maximum in healthy and moldy seeds of all groundnut varieties. Frequency of *A. flavus* was recorded in the range of 14.00 to 17.00 per cent on healthy and 18.00 to 26.00 per cent on moldy seeds of groundnut varieties. So seed health testing in groundnut is required to reduce the seed borne fungal pathogens.

ACKNOWLEDGEMENT

The authors are thankful to Dr. A. N. Sabalpara,

Directorate of Research and Dean P.G. Studies, NAU, Navsari for constant encouragement in presenting the information.

REFERENCES

- Anonymous 2012. District wise area, production and yield per hectare of important food and non-food crop in Gujarat state. Publ. Directorate of Agriculture, Gujarat State Krishi Bhavan, Paldi, Ahmedabad.
- Agrawal, S. C. and Gupta, R. K. 1989. Fungal pathogens detected on soybeans seeds grown in different localities. *Seed Research*, 17: 208-210
- Bhale, M.S., Khare, D., Rawat, N.D. and Singh, D. 2001. *Seed borne diseases objectionable in seed production and their management*. Scientific Publishers, Jodhpur (India). Pp: 10-16.
- Kushi, K. K. and Khare, M.N. 1979, Seed borne of pathogens of Sesame and their significance, *Seed Research* 7:48-53.
- Trividy, C.M. 1994 Loss of viability in storage microflora. *Seed Sci. and Technol.*, 1: 547 -562.